

Table D-1: Fish stockings in Weiss Lake, 1980 to 1994¹ (Source: Marion State Hatchery - ADCNR, as modified by Kleinschmidt).

Species	Date	Size Group (in.)	Total
Largemouth bass ²	1980	1-2	60,000
	1981	1-3	11,000
	1985	1-2	53,696
	1986	1-4	30,010
	1987	1-2	23,000
	1989	1-2	120,696
	1990	1	60,400
	1991	2-6	10,778
	1992	1-3	39,030
	1993	1	60,400
	1994	1	30,200
		Total	499,210
Striped bass ³	1985	1-2	29,700
	1987	1-2	30,000
	1988	1	21,560
			Total

¹In some years, the lake was stocked on more than one occasion, sometimes using different size classes. This table presents total number and size range stocked within a particular year.

²Florida strain (all largemouth bass stocked).

³Atlantic strain (all striped bass stocked).

Table D-2: Fish stockings in Henry Lake¹ (Source: Marion State Hatchery - ADCNR, as modified by Kleinschmidt)

Species	Date	Size Group (in.)	Total
largemouth bass ³	1981	1	18,860
	1984	1-3	25,800
	1986	1-2	30,032
	1987	1	11,400
	1988	2	22,470
	1989	1-2	36,300
	1990	1-2	10,555
	1991	1-2	22,446
	1992	1-2	22,240
	1993	1	22,240
		Total	222,343
striped bass ³	1976	1-2	11,000
	1977	1-2	57,025
	1983	1-2	58,530
	1984	1-2	55,000
	1985	2	55,000
	1995 ²	2	33,700
			Total
hybrid striped bass	1982	1	112,360
	1983	1-3	56,000
	1984	2	55,695
	1985	1-2	55,352
	1986	1-2	59,031
	1988	1-3	60,752
	1989	1-2	44,940
	1990	1-2	56,100
	1991	1-3	22,084
	1993	1	33,806
		Total	556,120

¹In some years, the lake was stocked on more than one occasion, sometimes using different size classes. This table presents total number and size range stocked within a particular year.

²Florida strain (all largemouth bass stocked)

³Gulf Coast strain

Table D-3: Fish stockings in Logan Martin Lake¹ (Source: Marion State Hatchery – ADCNR, as modified by Kleinschmidt)

Species	Date	Size Group (in.)	Total
largemouth bass	1964	2	7,800
	1981 ²	1-3	18,400
	1983	1-3	6,100
	1986	1-3	27,042
	1987	1-3	15,368
	1988	1-4	65,446
	1989	1	60,970
	1992	1-2	30,520
	1993	1	30,520
	1994	1-5	71,040
			Total 333,206
striped bass	1972	1-2	3,698
	1973	1-2	6,000
	1974	1-2	15,250
	1977	1-3	71,975
	1979	1-3	76,340
	1980	1	76,500
	1982	1-2	32,000
	1983	1-2	75,400
	1984	1-2	75,200
	1985	1-2	98,400
	1986 ³	1-2	70,800
	1987	1-4	92,000
	1988 ³	1-2	76,300
	1989	1-2	76,800
	1990	1-2	76,300
	1991 ³	1-3	76,616
	1992 ³	1-2	76,989
	1993 ³	1-2	80,798
	1994 ³	1	19,791
	1995 ³	1	76,351
1996 ³	1	76,255	
1998 ³	1	46,009	
1999 ³	1	12,345	
1999	1	33,932	
2001	1	40,120	
2002 ³	1	45,936	
2003 ³	2	76,567	
			Total 1,584,672

Table D-3: Fish stockings in Logan Martin Lake¹ (Source: Marion State Hatchery – ADCNR, as modified by Kleinschmidt)

Species	Date	Size Group (in.)	Total
hybrid striped bass	1976	1-2	35,451
	1977	1-3	3,640
	1981	1-2	37,000
	1982	1-3	154,770
	1983	1-3	157,462
	1984	1	157,820
	1985	1-2	157,014
	1986	1-2	157,100
	1987	1-2	94,500
	1988	1-2	45,800
	1989	2	30,000
	1990	1-2	30,500
	1991	1-3	30,823
	1992	1	30,520
	1993	1	30,745
	1994	1	45,916
	1995	1	76,585
	1996	1	61,310
	1998	1	47,464
	1999	1-2	45,873
2000	1	46,117	
2001	1-2	76,309	
2002	1	45,800	
2003	1-2	109,435	
			Total 1,707,954
Paddlefish	1981	22-29	46
	1983	3-4	3,500
			Total 3,546

¹In many years, the lake was stocked on more than one occasion, sometimes using different size classes. This table presents the total number and size range stocked within a particular year.

²Florida strain (all largemouth stocked since 1981)

³Gulf Coast strain (all other striped bass stocked were Atlantic strain)

Table D-4: Fish stockings in Lay Lake¹ (Source: Marion State Hatchery - ADCNR, as modified by Kleinschmidt)

Species	Date	Size Group (in.)	Total
largemouth bass ²	1983	1-2	38,650
	1986	1	20,000
	1987	1	12,800
	1988	1	52,300
	1989	1-3	55,396
	1990	2-4	6,458
	1991	1-2	24,000
	1992	1-2	24,000
	1993	1	27,363
	1994	1	24,000
	1995	1-5	83,535
	1996	2-4	71,595
	1997	1-4	267,310
	1998	1-4	249,522
	1999	1-4	191,868
	2000	1-3	182,105
			Total 1,330,902
striped bass	1975	1-3	59,710
	1977	1	24,150
	1978	1-4	50,390
	1979	1-2	42,600
	1982	1	68,119
	1983	1-2	66,480
	1984	1-3	60,300
	1985	1-2	74,400
	1986 ³	1-2	31,900
	1987	1-2	76,800
	1988 ³	1-2	54,200
	1989 ³	1-2	59,900
	1990 ³	1-2	60,100
	1991 ³	1-2	59,989
	1992 ³	1-2	70,968
	1993 ³	1-2	49,008
	1995 ³	1	60,070
	1996 ³	1	53,050
1997 ³	1	60,000	
1999	1	24,700	
2001	1	24,000	
2002 ³	1-2	24,000	
2003 ³	2	24,000	

Table D-4: Fish stockings in Lay Lake¹ (Source: Marion State Hatchery - ADCNR, as modified by Kleinschmidt)

Species	Date	Size Group (in.)	Total
			Total 1,178,834
hybrid striped bass	1976	1-2	61,176
	1982	1-3	60,000
	1983	1-2	60,000
	1984	1-3	99,800
	1985	2	100,019
	1986	1-3	101,600
	1987	1-2	70,800
	1988	1-2	36,000
	1989	1	36,100
	1990	1-2	36,000
	1991	1-2	36,000
	1992	1	36,062
	1993	1	36,098
	1994	1	36,078
	1995	1	36,000
	1996	1	31,400
	1997	1	36,313
	1998	1	25,390
	1999	1	36,020
	2000	1	24,808
	2001	1	24,000
	2002	1	24,000
	2003	1-2	24,480
			<hr/> Total 1,068,144

¹In many years, the lake was stocked on more than one occasion, sometimes using different size classes. This table presents the total number and size range stocked within a particular year.

²Florida strain (all largemouth stocked)

³Gulf Coast strain (unless noted, all other striped bass stocked were Atlantic strain)

Table D-5: Fish stockings in Mitchell Lake¹ (Source: Marion State Hatchery - ADCNR, as modified by Kleinschmidt)

Species	Date	Size Group (in.)	Total
largemouth bass ²	1983	1-2	13,300
	1986	1	12,000
	1987	1	6,100
	1988	1	23,400
	1990	1	11,875
	1991	2-3	11,800
	1992	1-3	45,682
	1993	1	11,700
	1994	1	11,700
		Total	147,557
striped bass ³	1975	1-2	29,250
	1980	1	29,000
	1984	1-2	30,100
		Total	88,350
hybrid striped bass	1982	2-4	65,616
	1983	1-3	29,000
	1984	2-3	29,500
	1985	1-2	60,600
	1986	1-2	60,600
	1988	1-2	29,500
	1989	1	17,000
	1990	1	23,400
	1991	1-2	23,400
	1992	1	12,524
	1993	1	97,580
	1994	1	23,400
	1995	1	23,400
	1996	1	48,600
	1997	1	23,400
	1998	1	23,369
	1999	1	23,433
2000	1	23,454	
2002	1	35,100	
2003	1-2	35,100	
		Total	707,976

¹In some years, the lake was stocked on more than one occasion, sometimes using different size classes. This table presents total number and size range stocked within a particular year.

²Florida strain (all largemouth bass stocked).

³Atlantic strain (all striped bass stocked).

Table D-6: Fish stockings in Jordan Lake¹ (Source: ADCNR – Marion State Hatchery, as modified by Kleinschmidt)

Species	Date	Size Group (in.)	Total
largemouth bass ²	1980	1-2	20,000
	1986	1	14,000
	1987	1	6,800
	1988	1-2	27,200
	1990	1	13,700
	1991	1-3	15,615
	1992	1-2	13,600
	1993	1	13,600
	1994	1	13,600
			Total
channel catfish	2003	3 – 9	153,142
striped bass ³	1975	1-2	69,000
	1976	1-2	60,495
	1977	1-2	62,550
	1978	1-2	62,000
	1979	1-4	70,070
	1980	1-2	23,800
			Total
hybrid striped bass	1981	1-3	63,580
	1982	1-4	89,020
	1983	1-3	69,002
	1984	1-2	69,300
	1985	1-3	69,000
	1986	1-2	69,000
	1987	1-2	40,300
	1988	1-2	33,200
	1989	1-2	27,200
	1990	1-2	27,200
	1991	2-3	26,378
	1992	1	27,200
	1993	1	67,500
	1994	2	27,200
	1995	1	27,207
	1996	1	43,354
	1997	1	27,200
	1998	1	27,386
	1999	1	26,857
2000	1	27,185	
2001	1	40,800	

Table D-6: Fish stockings in Jordan Lake¹ (Source: ADCNR – Marion State Hatchery, as modified by Kleinschmidt)

Species	Date	Size Group (in.)	Total
	2002	1	40,800
	2003	1-2	40,800
			Total 1,006,669

¹In many years, the lake was stocked on more than one occasion, sometimes using different size classes. This table presents the total number and size range stocked within a particular year.

²Florida strain (all largemouth stocked)

³Atlantic strain (all striped bass stocked)